



CHITTENANGO CENTRAL SCHOOLS
Michael R. Eiffe, Superintendent
1732 Fyler Road, Chittenango, NY 13037
Telephone: 315-687-2840—Fax: 315-687-2841

April 20, 2018

Jason P. Clark, Assistant
Superintendent for Instruction
315-687-2854-Fax: 315-687-2851

Scott Mahardy, Assistant
Superintendent for Business
315-687-2846-Fax: 315-687-2845

Mary Farber, Director
of Special Education/PPS
315-687-2844-Fax: 315-687-2851

Chittenango High School
Nicholas Fersch, Principal
Matthew Morkel, Asst. Principal
315-687-2900-Fax: 315-687-2924
Joseph Haven, Athletic Director
315-687-2905-Fax: 315-687-2924

Chittenango Middle School
Arnold Merola, Principal Grs. 5-6
Derek Sajnog, Principal Grs. 7-8
315-687-2800-Fax: 315-687-2801

Bolivar Road Elementary
Reneé Burgess, Principal
315-687-2880-Fax: 315-687-2881

Bridgeport Elementary
Jamie Durgey, Principal
315-687-2280-Fax: 315-687-2281

Food Services
Wendy Swift, Director
315-687-2866-Fax: 315-687-2861

Transportation
Connie Thorp, Supervisor
315-687-2870-Fax: 315-687-5823

Maintenance
Jeffrey Martin, Director
315-687-2863-Fax: 315-687-2861

Dear Parents/Guardians:

The Madison County Health Department has requested that parents of students attending Chittenango High School be informed of a confirmed case of pertussis (whooping cough) at the Chittenango High School. This correspondence and enclosure will provide additional information about pertussis and recommendations from the New York State Department of Health pertaining to individuals who might have been exposed. Pertussis is a contagious disease that is spread through the air by air droplets expelled with a cough or by direct contact with the droplets. Pertussis begins with mild upper respiratory cold-like symptoms, including a runny nose, low grade fever and a cough which becomes much worse over a one to two week period. Symptoms usually include a long series of coughs followed by a whooping noise. Older children, adults and very young infants may not develop the whooping noise. People with pertussis may have a series of coughs followed by vomiting, turning blue, or difficulty catching breath. The cough is often worse at night, and cough medicines usually do not help alleviate the cough.

If your child has been in contact with someone with pertussis, antibiotics may prevent him or her from becoming ill. If your child is already sick, giving antibiotics early can help your child get well faster and lower the chances of spreading the disease to others. You should watch for symptoms over the next several weeks.

Please consider the following New York State Department of Health recommendations, especially if you have younger children at home:

1. If your child comes down with cold symptoms that include a cough, talk to your child's doctor without delay. Tell the doctor that a confirmed pertussis case has occurred in your child's school.
2. Infants under one year old, especially those less than six months, are most likely to have severe symptoms if they develop pertussis. When possible, young infants should be kept away from people with a cough. **Infants with any coughing illness should be seen promptly by their doctor.**
3. Children under seven (7) years old that have not had at least five (5) doses of DTaP/DTP are at risk for getting the disease. Children between ages 11-18 years old who have not had a TdaP are also at risk for getting the disease. If you have children who have not been completely immunized against pertussis (particularly infants under one year), it is recommended you talk to your child's doctor about the benefits of vaccination.

The Tdap vaccine is a mandatory requirement for entrance into sixth grade. Parents may choose to get the vaccine sooner than sixth grade. If you have further concerns or questions, you may want to contact your doctor or the Madison County Health Department at 366-2848.

Sincerely,

Michael R. Eiffe
Superintendent of Schools

/kaq
Enclosure



Pertussis or Whooping Cough

What is pertussis?

Pertussis is also called whooping cough. It is a serious lung infection caused by bacteria. It is also very contagious and causes coughing fits. Whooping cough is most serious for babies with very small airways. It can lead to death.

What are the symptoms?

Symptoms usually appear about 7 to 10 days after exposure. But symptoms can appear from 4 to 21 days after a person is infected. It is rare but it can take up to 6 weeks to develop symptoms. Some people have milder symptoms, especially if they have been vaccinated.

Symptoms appear in 3 stages:

- Stage one lasts 1-2 weeks: runny nose, sneezing, low-grade fever, mild occasional cough. It may look like the common cold but the cough begins to get worse.
- Stage two can last 1-2 months: Cough becomes more severe. There are coughing fits that can be followed by a high-pitched whoop. The whoop means the person is trying to catch their breath. Sometimes a person will turn blue and vomit mucus or food. They also may have brief periods when they stop breathing.
- Stage three may last weeks to months: Recovery phase is when a person slowly gets better. Coughing fits can come back if the person gets another respiratory illness.

What are the complications of whooping cough?

Complications of whooping cough are more common in infants and young children. They may include pneumonia, middle ear infection, loss of appetite, sleep disturbance, fainting, dehydration, seizures, altered brain function (encephalopathy), brief periods when breathing stops and death.

How long is a person with whooping cough contagious?

A person with whooping cough can pass it to others as soon as they get cold-like symptoms. They can also pass it up to 3 weeks after they start coughing. If the infected person takes an appropriate antibiotic, they will not spread the germ after 5 full days of treatment.

What is the treatment?

Antibiotics can prevent and treat whooping cough. Recommended antibiotics include azithromycin, erythromycin and clarithromycin. Trimethoprim sulfamethoxazole can also be used.

If my child or another family member has been exposed to whooping cough, what should I do?

Call your local health department, doctor or clinic for advice. If you have close contact with an infected person, you will be given the same medicines used to treat whooping cough. This will help prevent you and others from getting sick.

What is the best way to prevent whooping cough?

The best way to prevent whooping cough is to get vaccinated.

We recommend that all children and adults get a pertussis-containing vaccine.

- The recommended pertussis vaccine for infants and children is called DTaP. This protects children against 3 diseases: diphtheria, tetanus, and whooping cough (pertussis).

- DTaP shots are given at 2, 4, and 6 months of age. To maintain protection, boosters are also given at 15 through 18 months and 4 through 6 years of age.
- 7- to 10-year-olds who are not fully immunized with DTaP, should get a Tdap vaccine which protects against diphtheria, tetanus and whooping cough (pertussis).
- Preteens going to the doctor for their regular checkup at 11 or 12 years of age should get a booster dose of Tdap. If a preteen did not get this Tdap at 11 to 12 years, they should be vaccinated at their next doctor visit.
- All adults 19 and older who have not received a Tdap vaccine, need a booster shot.
- Adults should receive a tetanus and diphtheria (Td) booster every 10 years. They should also substitute a Tdap vaccine for one of the boosters. The Tdap can be given earlier than the 10-year spacing.
- It is very important for adults to make sure they are up-to-date with their pertussis-containing vaccines if they spend any time with infants. This includes parents, grandparents, siblings, babysitters, relatives and friends.
- Pregnant women should receive Tdap in the third trimester of every pregnancy to pass immunity to their newborn until the baby is old enough to begin shots at 2 months of age.
- Travelers should be up-to-date with pertussis-containing vaccines before they travel. Whooping cough occurs everywhere in the world.
- Health care staff who have direct contact with patients should get a single dose of Tdap if they have not already received it.

What are the whooping cough vaccine requirements for school attendance?

- Prekindergarten (Day Care, Head Start, or Nursery): 4 doses (age-appropriate)
- Kindergarten through 5th grade: 5 doses, or 4 doses with the 4th dose at 4 years of age or older. Or, 3 doses if they start the series at 7 years of age or older
- Grades 6 through 12th grade: 3 doses
- Grades 6 to 12: one dose of Tdap (required for students enrolling in grades 6 to 12 who have not previously received Tdap at 7 years of age or older)

What should I do if I'm not sure I received whooping cough vaccine?

Check with your doctor. An adolescent or adult without documentation of having received a primary series of at least three doses of tetanus and diphtheria (Td) vaccines in their lifetime should receive this series. The first dose should be Tdap. The remaining two doses should be adult formulation Td.

What should I or my family do to prevent whooping cough if we travel out of the country?

Whooping cough is still common in the United States and in many other countries. Make sure that you and your children are fully vaccinated before traveling.

- Adults 19 or older should receive a single dose of Tdap vaccine. Confirm with your doctor that you have received the vaccine for pertussis.
- Infants, children and adolescents should be up-to-date with pertussis-containing vaccines as described above.

Travel and whooping cough:

wwwnc.cdc.gov/travel/diseases/pertussis

How can I find out about whooping cough outbreaks?

www.cdc.gov/pertussis/outbreaks/about.html

Learn more about whooping cough:

www.cdc.gov/pertussis

For more information about vaccine-preventable diseases: www.health.ny.gov/prevention/immunization/

Bureau of Immunization